

Frequently Asked Questions about Bovine Tuberculosis

The diagnosis of four herds infected with bovine Tuberculosis in the first 5 months of 2013 is an reminder that all Michigan dairy and beef cattle producers should understand what they can do to reduce the risk of disease in their herd.

Posted on **May 31, 2013** by **Phil Durst** (http://msue.anr.msu.edu/experts/phillip_durst), Michigan State University Extension

What causes Bovine Tuberculosis or TB?

TB is caused by an infection with the bacteria called *Mycobacterium bovis* (M. bovis). Infected animals may not show any outward signs of disease until the most advanced stages.

Can M. bovis survive in the environment?

M. bovis is very resilient. Survival is affected greatly by temperature and moisture. When exposed to summer temperatures, sunlight and drying conditions, the bacteria dies, but away from sunlight and in moist conditions in soil and manure, M. bovis can survive for many months, especially in the cold.

How is the disease transmitted from wildlife?

In Michigan, as well as in areas of New Zealand, the U.K. and other locations, wild animals are believed to be a primary major source of M. bovis transmission to cattle. A wildlife reservoir of the disease occurs because of three factors: 1. The disease has established itself in a wildlife population, 2. The live bacteria can be shed by the animal, and 3. There is opportunity for M. bovis to be transmitted to cattle through both direct and indirect pathways.

An example of direct transmission of M. bovis would be when aerosols or droplets that are exhaled or coughed from an infected animal and contain the bacteria go directly from infected animal to a susceptible animal. This type of transmission would most likely occur in crowded conditions or high populations or in contained areas.

Indirect transmission could occur when bacteria from an infected animal is transmitted to a susceptible animal through some indirect means. Examples would be when an infected animal's respiratory secretion or saliva containing M. bovis is deposited on a feedstuff which is then subsequently ingested by a susceptible animal, or through ingesting milk that contains the live M. bovis.

In Michigan, it is believed that indirect transmission of M. bovis is the most common cause of herd infections.

Can *M. bovis* be spread through small wild animals like opossums and raccoons?

While the disease has been identified in these animals as well as in scavenging carnivores such as coyotes, bobcat, fox and bear, it is not believed that these animals can pass it on to cattle or that it has become endemic in their populations. Producers are encouraged to control small animals on their farms to suppress possible transmission of various diseases.

How contagious is *M. bovis* within a cattle herd?

Diseases spread when conditions favor exposure and when animals are susceptible. In Michigan herds located in the geographic area known to have TB in wildlife, annual testing of cattle has found the infection often in early stages. In most of the infected herds, only one or two infected animals have been found. From that, it was believed that the disease was not very contagious. However, the experience with a few herds in which a high proportion of animals has been found infected causes us to question that conclusion. In fact, when conditions and management result in greater exposure, particularly of young animals, the disease can apparently be fairly contagious.

Can cattle get TB from feed?

Yes, cattle as well as deer and other animals that eat feed on which saliva from infected animals has been deposited, and where the bacteria survives, can get infected. That has been shown in research in which infected animals were allowed to eat at feed and then were removed and that feed was fed to uninfected animals, all of which became infected. That is the scientific basis for eliminating the feeding and baiting of wild deer that can spread the disease within their herds.

In fact, survival of *M. bovis* on common deer bait feeds such as apples, carrots, corn, sugar beets and potatoes in an experimental study lasted at least 7 days on all materials and could still be isolated from apples, corn and beets at 112 days. It is believed that cattle grazing on pasture and eating apples from trees in pastures where deer have had access can be a means of transmission.

However, the question of whether harvested feed, such as hay, is a source of transmission has greater uncertainty. Experimentally, when dry hay was inoculated with *M. bovis*, survival in fall and winter was weeks or even months, however, survival in summer was less than 3 days. If saliva with bacteria is deposited on forage that is cut for hay or haylage, and cured in the sun for hours or days, there is lower likelihood of its survival. With haylage the process of ensiling results in the production of acids that preserve the feed. The impact of the ensiling process on the survival of *M. bovis* is under investigation in a 2013 study at Michigan State University. The hypothesis is that the bacteria cannot survive long and the intent is to characterize the death curve of the bacteria over time.

What other animals on my farm could get bovine TB?

Horses and sheep are apparently resistant to TB and are not a concern. Goats and pigs can become infected. In addition, dogs and cats that drink milk from infected cows or eat tissues from infected cattle may also become infected.

Can people get bovine TB?

Though the disease is called bovine Tuberculosis, people indeed may become infected with it, however, it is not a very high risk. Simply being around infected cattle and working with infected cattle has not resulted in human infections. The most likely way that humans would become infected is by drinking unpasteurized milk from a herd with infected cows. It was that means of transmission that provided the impetus years ago for laws requiring the pasteurization of milk for sale.

Can't TB be treated?

Infected people can be treated for TB and in the vast majority of cases the treatment is successful. The course of treatment takes 6 to 9 months and is, therefore, expensive. Treatment of infected cattle is not feasible.

How safe is our food?

Pasteurization kills disease pathogens meaning that pasteurized milk is completely safe for all consumers. The USDA meat inspection system examines carcasses and looks for any type of problem that would impact the health of consumers. When combined with proper handling and cooking, our meat supply is also safe for all consumers.

How reliable is the TB test?

Tests are evaluated for their sensitivity (having a low rate of false negative responses) and their specificity (having a low rate of false positive responses). In diagnosing TB, no test is perfect. A test that has high sensitivity, like the caudal fold test (CFT), is used as a screening test, and is followed up with a test that has a higher specificity (gamma interferon or comparative cervical test, CCT). When an animal responds on both of these tests, it is taken and sacrificed and the tissues and lymph nodes examined. Samples from these will be tested for genetic markers (PCR) of *M. bovis* and cultures will be done to see if lab personnel can grow the bacteria from tissue samples. If either of these last two tests is positive, the animal is called infected, and therefore the herd is called infected.

CFT has been the standard screening test for many years. Its sensitivity is only around 80 percent meaning that approximately 20 percent of infected animals will go undetected. The specificity of the CFT is about 95 percent meaning that 2-5 percent of the time it will yield a false positive result. The desire and effort within the scientific community has been to have a screening test with a higher sensitivity and higher specificity and research in developing other

tests is ongoing.

If I'm a cattle producer what should I do?

[Michigan State University Extension \(/\)](#) recommends taking the following measures to reduce the risk of transmission of TB to your herd:

- **Wildlife control.** Wildlife can be vectors of diseases, that is, they can carry diseases to cattle. This is not just the case with TB but also with other diseases as well. Cattle producers, no matter where they are located, should develop measures to protect feed, water and housing areas from wildlife and should use all legal means to control the numbers of wild animals in the area of their farm. Talk with your Department of Natural Resources Officer to learn more about the legal means available to you.
- **Herd health.** Cattle producers need to support the health of their herd through every means possible and lower the potential exposure to all disease pathogens because one disease may impair immune response to another disease, making them more susceptible. This includes having a clean and dry environment for all cattle, particularly at calving, managing the stocking rate to keep aerosol and manure exposure lower, reducing the mixing of cattle of different ages and from different herds, and reducing the exposure of cattle to blood from other cattle. The latter may mean that you and your veterinarian commit to single use of examination sleeves and needles, and reduced use of bulls. In addition cattle should be fed diets that meet their nutritional and immune system needs including proper balancing for minerals and vitamins. Additionally, vaccination is an important tool to strengthen the immune response to certain viruses and bacteria. An animal whose health is compromised by one disease is more susceptible to other diseases.
- **Work with your veterinarian.** Managing cattle herd health should always be a partnership with your veterinarian in which together you evaluate the weaknesses and breakdowns in herd health and together plan ways to strengthen your herd health program.
- **Always properly identify animals.** Whenever an infected animal is found, the first question is what other animals have been exposed. Tracing the movements of an animal and being able to determine other exposed animals will help control contagious diseases faster with fewer animals needing to be killed. That requires that all animals that move be identified and that good records be kept of animals and where they went or where they came from. Therefore, it goes beyond simply complying with regulations to having records that can be provided if and when necessary.

This article was published by [Michigan State University Extension \(http://www.msue.msu.edu\)](http://www.msue.msu.edu). For more information, visit [http://www.msue.msu.edu \(http://www.msue.msu.edu\)](http://www.msue.msu.edu). To have a digest of information delivered straight to your email inbox, visit <http://bit.ly/MSUENews> (<http://bit.ly/MSUENews>). To contact an expert in your area, visit <http://expert.msue.msu.edu> (<http://expert.msue.msu.edu>), or call 888-MSUE4MI (888-678-3464).

Related Articles

[Proper manure management is important for everyone, including small-scale livestock or horse farms](http://msue.anr.msu.edu/news/proper_manure_management_is_important_for_everyone_including_small-scale_livestock_or_horse_farms)
([http://msue.anr.msu.edu/news/proper manure management is important for ever](http://msue.anr.msu.edu/news/proper_manure_management_is_important_for_ever)

[yone including small scale li](#)

August 19, 2014 | **Shelby Bollwahn** | Using best management practices can help protect the environment. These practices can also improve the health and well-being of your animals and increase your farm's profit.

[Corn is again a low cost feedstock for biofuel](#)

[http://msue.anr.msu.edu/news/corn is again a low cost feedstock for biofuel](http://msue.anr.msu.edu/news/corn_is_again_a_low_cost_feedstock_for_biofuel)

August 6, 2014 | **Mark Seamon** | Corn prices are less than half of the price at this time in 2013.

[Timing is right for summer alfalfa seeding](#)

[http://msue.anr.msu.edu/news/timing is right for summer alfalfa seeding](http://msue.anr.msu.edu/news/timing_is_right_for_summer_alfalfa_seeding)

August 5, 2014 | **Phil Kaatz** | Long-term effects of winterkill in alfalfa fields include reduced yields and shorter stand life.

[Michigan Crude Oil And Petroleum Act 16 of 1929 and landowners: Part 3](#)

[http://msue.anr.msu.edu/news/the michigan crude oil and petroleum act 16 of 1929 and landowners part 3](http://msue.anr.msu.edu/news/the_michigan_crude_oil_and_petroleum_act_16_of_1929_and_landowners_part_3)

August 1, 2014 | **Curtis Talley Jr.** | As energy infrastructure for our state and nation improve and grow, a landowner may be approached to allow the construction of a pipeline across their property.

[Certified Animal Welfare Audits for Farms: Part 6 - Follow-up for certification](#)

[http://msue.anr.msu.edu/news/certified farm animal welfare audits for farms part 6 follow up for certifi](http://msue.anr.msu.edu/news/certified_farm_animal_welfare_audits_for_farms_part_6_follow_up_for_certifi)

July 28, 2014 | **Tina Conklin** | The final installment in the six part series of Certified Farm Animal Welfare Audits looks at the follow-up corrective actions that need to take place after the audit in order for your farm to achieve certification.

Related Events

[Smart Drainage Field Day: Use It, Don't Lose It!](#)

[http://msue.anr.msu.edu/events/smart drainage field day use it dont lose it](http://msue.anr.msu.edu/events/smart_drainage_field_day_use_it_dont_lose_it)

Date: Aug 28, 2014

Location: 2435 N. Tallman Road Fowler, MI

[Grazing School \(\[http://msue.anr.msu.edu/events/grazing school\]\(http://msue.anr.msu.edu/events/grazing_school\)\)](#)

Date: Sep 9, 2014 - Sep 10, 2014

Location: MSU Kellogg Biological Station Pasture Dairy Center, 10461 N 40th, St., Hickory Corners, MI 49060

[4-H Veterinary Science Teen and Adult Leaders Workshop](#)

[http://msue.anr.msu.edu/events/4 h veterinary science teen and adult leaders workshop](http://msue.anr.msu.edu/events/4_h_veterinary_science_teen_and_adult_leaders_workshop)

Date: Feb 27, 2015 - Feb 28, 2015

Location: Kettunen Center, 14901 4-H Drive, Tustin, MI 49688

